

EFFICACY AND DURABILITY OF FOSTEMSAVIR, IBALIZUMAB, OR LENACAPAVIR-INCLUDING REGIMENS IN PEOPLE LIVING WITH MULTIDRUG-RESISTANT HIV-1: RESULTS FROM THE PRESTIGIO REGISTRY

ePOSTER number:eP.B1.031

G. Torkjazi^{1,2}, L. Galli³, R. Gagliardini⁴, M. Feasi⁵, S. Bonora⁶, F. Lagi⁷, E. Focà⁸, G. Marchetti⁹, A. Cervo¹⁰, F. Maggiolo¹¹, R. Gulminetti¹², F. Vichi¹³, R. Papaioannu Borjesson¹, A. Castagna^{1,3}, V. Spagnuolo³, on behalf of the PRESTIGIO Study Group

1. Vita-Salute San Raffaele University, School of Medicine, Milan, Italy; 2. Università La Sapienza, Facoltà di Medicina e Chirurgia, Rome, Italy; 3. IRCCS Ospedale San Raffaele, Unit of Infectious Diseases, Milan, Italy; 4. National Institute for Infectious Diseases 'L. Spallanzani' IRCCS, Rome, Italy; 5. Ente Ospedaliero Ospedali Galliera, Genoa, Italy; 6. University of Turin, Department of Medical Sciences, Turin, Italy; 7. University of Florence, Department of Clinical and Experimental Medicine, Florence, Italy; 8. ASST Spedali Civili Hospital, University of Brescia, Division of Infectious and Tropical Diseases, Brescia, Italy; 9. San Paolo Hospital, ASST Santi Paolo e Carlo, Department of Health Sciences, University of Milan, Clinic of Infectious Diseases and Tropical Medicine, Milano, Italy; 10. Azienda Ospedaliero-Universitaria di Modena, Department of Infectious Diseases, Modena, Italy; 11. Azienda Ospedaliera Papa Giovanni XXIII, Department of Infectious Diseases, Bergamo, Italy; 12. Fondazione IRCCS Policlinico San Matteo, Department of Medical Sciences and Infectious Diseases, Pavia, Italy; 13. Santa Maria Annunziata Hospital, Infectious Diseases Unit, Bagno a Ripoli, Italy

19th EUROPEAN AIDS CONFERENCE
18-21 October, 2023 | Warsaw, Poland

BACKGROUND AND AIMS

Heavily treatment-experienced people living with HIV (HTE-PLWH) often have few, if any, treatment options due to archived drug resistance mutations and/or intolerance to therapy^{1,2}. Therefore, the availability of drugs with novel mechanisms of action in this population is essential to achieve viral suppression, restore immune function and prevent clinical progression³. The aim of this study was therefore to describe the efficacy and durability of fostemsavir (FTR), ibalizumab (IBA) or lenacapavir (LEN) containing regimens in people living with 4-drug class resistant HIV (4DR-PLWH) in a real-life setting.

STUDY DESIGN AND METHODS

We included 4DR-PLWH from the PRESTIGIO registry (NCT04098315) treated with a regimen containing FTR, IBA or LEN and with documented resistance to NRTIs, NNRTIs, PIs or INSTIs. FTR, IBA or LEN discontinuation was defined as interruption of the drug for any reason. Follow-up (FU) started from the date of initiation of FTR, LEN or IBA (baseline; BL) until discontinuation of FTR, IBA, or LEN or death/freezing date (30 June 2023).

Descriptions are reported as median (IQR) or frequency (%). The genotypic susceptibility score for the optimized background therapy introduced with FTR, LEN, or IBA was estimated according to the cumulative data of the available plasma genotyping resistance tests recorded for each individual.

RESULTS

Among 35 4DR-PLWH, we considered 27, 11 and 10 FTR-, IBA- and LEN-including regimens, respectively (Table 1).

FTR-including regimens: at the end of FU [median duration 18.7 (5.6-82.7) months], virological efficacy (VE; HIV-RNA <200 copies/mL) was 66.7% (18/27); 8/27 (29.6%) regimens had been discontinued. Among people still in a FTR-including regimen, the median change in CD4+ was +97 (-38/+182) cells/mm³ (p=0.064).

IBA-including regimens: median FU 14.1 (5.6 -39.1) months, VE was 63.6% (7/11); 4/11 (36%) regimens had been discontinued. Median CD4+ change, in people still on treatment with IBA, was +63 (-38/+122) cells/mm³ (p=0.109).

LEN including regimens: median FU 30.3 (18.7-33.5) months, VE was 90% (9/10); 2/10 (20%) regimens had been discontinued. Median CD4+ change , in people still on treatment with LEN, was +5 (-48/+101) cells/mm³ (p=0.742).

Regimens including a concomitant administration of FTR, IBA and LEN (n=4) are described in Table 2.

CONCLUSIONS

- In 4DR-PLWH, the overall efficacy and durability of FTR, IBA or LEN-including regimens were good.
- Efficacy and safety data on combination regimens including entry and capsid inhibitors are urgently needed in this fragile population.

Table 1: Characteristics of PWH at baseline (Panel A) and at the end of follow-up (Panel B).

Panel A

Characteristic	Category	FTR-including regimens (n=27)	IBA-including regimens (n=11)	LEN-including regimens (n=10)
Age (years) at baseline		54.3 (47.6 - 58.6)	55.1 (31.7 - 58.2)	52.1 (45.4 - 65.3)
Years of HIV infection at baseline		26.6 (21.4 - 33.1)	28.8 (25.9 - 33.3)	28.8 (22.8 - 30.7)
Years of ART at baseline		23.2 (19.2 - 25.9)	26.5 (23.9 - 31.6)	25.4 (22.6 - 28.7)
Viral Subtype				
	B	21 (78.3%)	8 (72.7%)	8 (80%)
	Non-B	6 (21.7%)	3 (27.2%)	2 (20%)
R5 tropism at baseline		11 (41%)	5 (45%)	4 (40%)
Nadir CD4+ (cells/mm ³)		144.5 (29 - 204)	29 (5 - 137)	40.5 (3 - 137)
Baseline HIV-RNA (log ₁₀ copies/ml)		3.86 (2.09 - 4.91)	4.43 (3.47 - 5.06)	3.46 (3.05 - 4.33)
Baseline CD4+ (cells/mm ³)		244 (141 - 641)	167 (9 - 324)	287 (173 - 593)
OBT GSS		1.5 (1 - 2)	1.5 (1 - 2.5)	2 (1 - 2.5)

Panel B

Characteristic	Category	FTR-including regimens (n=27)	IBA-including regimens (n=11)	LEN-including regimens (n=10)
HIV-RNA at last visit (copies/mL)				
	<50	13 (48.2%)	5 (45.5%)	7 (70%)
	≥50 - <200	5 (18.5%)	2 (18.2%)	2 (20%)
	≥200	9 (33.3%)	4 (36.3%)	1 (10%)
CD4+ at last visit (cells/mm ³)		328.5 (208 - 616)	131 (105 - 230)	282 (159 - 630)
Change in CD4+ during FU (cells/mm ³)	Among those still on treatment	73 (-27/+179)	53 (-13/91)	-10 (-48/+52.5)
Duration of FTR, IBA or LEN-regimen (months)		18.7 (5.6 - 82.7)	14.1 (5.6 - 39.1)	30.3 (18.7 - 33.5)
Discontinuation of FTR, IBA or LEN-regimen		8 (29.6%)	4 (36.4%)	2 (20%)
Reasons of discontinuation				
	Virological failure	5 (63%)	1 (25%)	0
	Toxicities/Other	3 (37%)	3 (75%)	2 (100%)

OBT-GSS: optimized background therapy-genotypic susceptibility score; FTR: fostemsavir; IBA: ibalizumab; LEN: lenacapavir

Table 2: Regimens including a concomitant administration of fostemsavir (FTR), ibalizumab (IBA) or lenacapavir (LEN)

ID PLWH	New drugs Combination	Optimized background treatment (OBT)	Baseline Cumulative Genotype	Date of start	Date of stop	OBT GSS	BL HIV-RNA (copies/mL)	Last HIV-RNA (copies/mL)	BL CD4+ (cells/μL)	Last CD4+ (cells/μL)	Duration (months)	Discontinuation (reason)
104-002	IBA+LEN	FTC, TAF, DOR, DRV/r, MVC	RT:41L,74V,138A,181CIV,184V,190A,210W,215F,215Y,219E,230L PRO:32I,33F,46I,46L,54A,54L,54T,54V,82F,84V,90M IN: 140S,148H	26 Oct 2020	17 May 2022	1.0	26800	<1	42	159	18.7	Yes, only LEN: patient's decision
108-0021	IBA+FTR+LEN	3TC, TDF, DOR, MVC	RT: 67N,70R,138K, 181C, 184V,190A, 215F,215Y,219Q PRO: 32I,46I,76V,82A,84V IN: 138T,140S,143C,148H	18 Jan 2021	ongoing	1.5	2982	52	734	721	28.7	No
120-001	IBA+FTR	FTC, TAF, DRV/r, DTG, MVC	RT:67N,70R,181C, 184V,190A, 219Q PRO:47V,50V,54V IN:140S,148H	22 Dec 2022	ongoing	2.5	396	23	324	286	5.6	No
134-005	IBA+FTR+LEN	FTC, TAF, DRV/r, DTG	RT:41L,74V,103N,181C, 184V,215Y PRO:33F,84V,90M IN:138K,140S,148H	12 Sep 2022	ongoing	1.0	2400	20	296	230	8.9	No

GSS: susceptibility score; RT: reverse transcriptase; PRO: protease; IN: integrase

Acknowledgments

PRESTIGIO Study Group: STEERING COMMITTEE: Antonella Castagna (Coordinator), Vincenzo Spagnuolo, Laura Galli, Franco Maggiolo, Leonardo Calza, Emanuele Focà, Giovanni Cenderello, Antonio Di Biagio, Giulia Marchetti, Stefano Rusconi, Adriana Cervo, Roberta Cagliardini, Stefano Bonora, Anna Maria Cattelan, Maurizio Zazzi, Maria Mercedes Santoro; VIROLOGY TEAM AND BIOLOGICAL BANK: Maurizio Zazzi, Maria Mercedes Santoro, Andrea Galli, Francesco Saladini, Daniele Armenia; STUDY COORDINATORS: Elisabetta Carini, Sabrina Bagaglio; STATISTICAL AND MONITORING TEAM: Laura Galli, Riccardo Lolatto, Sara Diotallevi; ENROLLING CENTERS: ANCONA: Marcello Tavio, Alessandra Mataloni Paggi; BARI: Annalisa Sarcino, Flavia Balena; BERGAMO: Franco Maggiolo, Laura Corni, Daniela Valenti, Claudia Suardi; BOLOGNA: Leonardo Calza, Malerba Federica; BRESCIA: Francesco Castelli, Emanuele Focà, Davide Minisci, Francesca Pennati, Anna Celotti, Francesca Brognoli; BUSTO ARSIZIO: Barbara Menzaghi, Maddalena Farinazzo; CATANIA: Bruno Cacopardo, Maurizio Celestia, Michele Salvatore Paternò Raddusa, Carmen Giarratana; CATANZARO: Carlo Torti, Paolo Fusco, Gabriele Bruno; CREMONA: Angelo Pan, Paola Brambilla, Chiara Fornabai; FIRENZE: Alessandro Bartoloni, Filippo Lagi, Susanna Giache, Francesca Vichi, Francesco Maria Fusco, Alessio Bellucci, Elisa Mirabelli, Paola Corsi, Seble Tekle Kiro, Filippo Ducci; FOGGIA: Teresa Santantonio, Sergio Lo Caputo, Sergio Ferrara, Marianna Narducci; GENOVA: Emanuele Pontali, Marcello Feasi, Antonio Sara, Matteo Bassetti, Antonio Di Biagio, Sabrina Bianchi; MILANO: Antonella Castagna, Vincenzo Spagnuolo, Elisabetta Carini, Sabrina Bagaglio, Laura Galli, Riccardo Lolatto, Andrea Galli, Rebecca Papaioannu, Tommaso Clemente, Sara Diotallevi, Spinelio Antinori, Tiziana Formenti, Giulia Giacomelli, Giulia Marchetti, Lidia Gazzola, Federica De Flavis, Massimo Puoti, Cristina Molit, Federico D'Amico; MODENA: Cristina Mussini, Adriana Cervo, Enrica Roncaglia, Giulia Nardini, Barbara Baghetto; NAPOLI: Elia Mancillo, Amedeo Lanzardo; PADOVA: Anna Maria Cattelan, Maria Mazzitelli; PALERMO: Antonio Cascio, Marcello Trizzino; PARMA: Elisa Fronti, Diletta Lacabue; PAVIA: Roberto Gulminetti, Andrea Zuccarini; PERUGIA: Daniela Francis, Elisabetta Schiaroli, Giuseppe De Socio; REGGIO EMILIA: Elisa Garlassi, Romina Corsini; ROMA: Roberta Cagliardini, Marisa Fusto, Loredana Sarmati, Vincenzo Malagolino, Tiziana Mulas, Simona Di Giambenedetto, Silvia Lamonic; SANREMO: Giovanni Cenderello, Rachele Pincino; SIENA: Mario Tumbarello, Massimiliano Fabiani, Francesca Panza, Iaria Rancan; TORINO: Giovanni Di Perri, Stefano Bonora, Micol Ferrara; VERONA: Marina Malena, Marta Fiscon. SUPPORTED BY: VIV Healthcare, Gilead Sciences, Theratechnologies, MSD

References

1. EACS Guidelines. Version 11.1 October 2022. Accessed September 30, 2023. https://www.eacsociety.org/media/guidelines-11.1_final_09-10.pdf
2. Galli L, Parisi MR, Poli A et al. Burden of Disease in PWH Harboring a Multidrug-Resistant Virus: Data From the PRESTIGIO Registry. Open Forum Infect Dis. 2020;7:ofaa456.
3. Mucchi C, Canetti D, Castagna A, et al. Efficacy and Safety Profile of Fostemsavir for the Treatment of People with Human Immunodeficiency Virus-1 (HIV-1): Current Evidence and Place in Therapy. Drug Des Devel Ther. 2022;16:297-304.

Contact Information

Vincenzo Spagnuolo, MD.
Infectious Diseases Unit San Raffaele Scientific Institute, Milan, Italy,
telephone: +390226437907, e-mail: spagnuolo.vincenzo@hsr.it